

EXPLORING THE SURFACE ROUGHNESS OF SMALL ICE CRYSTALS BY MEASURING HIGH RESOLUTION ANGULAR SCATTERING PATTERNS

MARTIN SCHNAITER,^{a*} PAUL HENRY KAYE,^b EDWIN HIRST,^b
ZBIGNIEW ULANOWSKI,^b AND ROBERT WAGNER^a

ABSTRACT. Surface roughness of atmospheric ice particles is an important yet poorly investigated microphysical property in the context of the climate impact of cirrus and mixed-phase clouds. Measurements of single particle two-dimensional light scattering patterns have been shown to be a promising method to identify particle roughness in natural clouds. This method was applied in a laboratory study on the surface roughness properties of small ice particles that have been grown under simulated cirrus conditions.

1. Introduction

The microphysical properties of ice crystals play an important role for the radiative impact of cirrus and mixed-phase clouds as they determine the angular single scattering phase function of the particles and in this way the shortwave bulk optical properties of the clouds [1, 2]. In order to compile representative parameterisations of single scattering properties of natural ice particles, detailed *in situ* microphysical and optical measurements are therefore required which are also prerequisites for the evaluation of remote sensing measurements. Such *in situ* investigations are mostly restricted to particles larger than about 50 μm , mainly because of optical resolution limitations and potential impacts by ice crystal breakup on the probe inlets. Furthermore, the ice particle surface roughness is by far the least investigated microphysical property addressed by *in situ* measurements and is mainly limited to indirect evidence like the shape of the scattering function [3, 4].

The use of high resolution measurements of the two-dimensional (2D) single particle scattering pattern is a promising way to close these gaps [5]. The latest version of the Small Ice Detector, named SID-3, which has been developed at the University of Hertfordshire is the first *in situ* cloud probe that measures highly resolved 2D scattering patterns. First *in situ* SID-3 scattering patterns of atmospheric ice particles have been analysed to identify surface roughness using statistical measures of the Grey-Level Co-occurrence Matrix




Figure 1. Time evolution of the AIDA ice cloud characterisation experiment HALO06 #47. Expansion cooling was started at $t=0$ sec and stopped at $t=570$ sec. The nucleation of ice by deposition freezing on the surface of the pre-existing soot particles was initiated at $t=73$ sec when the relative humidity with respect to ice had been reached a saturation ratio of 125%. The ice cloud formation and evaporation is clearly reflected by the time evolution of the scattering and depolarisation measurements.

(GLCM) [6]. Here, we examine the GLCM analysis to extract surface roughness parameters from 2D scattering patterns in case of small ~ 10 micrometer-sized ice crystals that have been generated under controlled conditions in the laboratory.

2. Results

Ice cloud characterisation experiments were conducted at the large cloud simulation chamber AIDA of the Karlsruhe Institute of Technology [7]. Figure 1 shows the temporal evolution of the thermodynamic conditions of the cloud expansion experiment discussed herein. The experiment was conducted at an initial temperature of 229 K and with submicron soot particles as nuclei for the ice crystal formation. During the expansion cooling of the chamber volume the relative humidity with respect to ice increased to about 125 % before the ice cloud started to form. The growing ice particles reduced the supersaturation and confined the relative humidity to ice saturated conditions before the expansion was stopped at 570 s experiment time. Corresponding results of SID-3 measurements are shown in Figure 2. The comparison of particle sizes deduced from the intensity of the individual 2D scattering patterns and retrieved from FTIR extinction spectroscopy of the ice cloud shows good agreement and reveals a mean equivalent particle size of about $10 \mu\text{m}$ when the cloud has been fully evolved. The mean azimuthal intensity profile from each




Figure 2. Analysis of the single particle 2D scattering patterns acquired by the SID-3 probe during the AIDA cloud experiment HALO06 #47. Upper panel: The size and shape analysis of the cloud particles reveals a dominating presence of small columnar ice crystals with a mean volume equivalent diameter of about $10 \mu\text{m}$. The GLCM texture feature energy - which is the sum of squared elements of the GLCM - shows a useful discrimination according to the degree of surface roughness of the columnar ice crystals. Lower panel: Representative 2D patterns of the four GLCM energy ranges >0.6 (a), $0.5-0.55$ (b), $0.4-0.45$ (c), and $0.34-0.37$ (d), indicated as shaded areas in the upper panel. Each individual image covers an angular range from 7° to 24° .

individual 2D scattering pattern image was calculated and Fourier transformed in order to get a shape discrimination, i.e. selecting columnar particles [8]. We investigated five statistical features of the GLCM – entropy, dissimilarity, contrast, homogeneity, and correlation

– for their surface roughness discrimination capability for this particle subset. Energy was found to be a very robust measure of particle roughness with the potential to effectively discriminate particles with different degrees of roughness.

Acknowledgments

This work was supported by the German Research Foundation under the grant SCHN 1140/1-1 as part of the priority program 1294.

References

- [1] A. J. Baran, “A review of the light scattering properties of cirrus”, *Journal of Quantitative Spectroscopy and Radiative Transfer* **110**, 1239 – 1260 (2009)
- [2] A. Ehrlich, M. Wendisch, E. Bierwirth, A. Herber, A. Schwarzenböck, “Ice crystal shape effects on solar radiative properties of Arctic mixed-phase clouds–Dependence on microphysical properties”, *Atmospheric Research* **88**, 266 – 276 (2008)
- [3] V. Shcherbakov, J.-F. Gayet, B. Baker, P. Lawson, “Light Scattering by Single Natural Ice Crystals”, *Journal of the Atmospheric Sciences* **63**, 1513 – 1525 (2006)
- [4] O. Jourdan, G. Mioche, T. J. Garrett, A. Schwarzenböck, J. Vidot, Y. Xie, V. Shcherbakov, P. Yang, J.-F. Gayet, “Coupling of the microphysical and optical properties of an Arctic nimbostratus cloud during the ASTAR 2004 experiment: Implications for light-scattering modeling”, *Journal of Geophysical Research* **115**, D23206, doi:10.1029/2010JD014016 (2010)
- [5] P. H. Kaye, E. Hirst, R. S. Greenaway, Z. Ulanowski, E. Hesse, P. J. DeMott, C. Saunders, P. Connolly, “Classifying atmospheric ice crystals by spatial light scattering”, *Optics Letters* **33**, 1545 – 1547 (2008)
- [6] Z. Ulanowski, E. Hirst, P. H. Kaye, R. S. Greenaway, “Light scattering by ice particles in the Earth’s atmosphere and related laboratory measurements”, in *12th Int. Conference on Electromagnetic & Light Scattering*; Helsinki, Finland (2010), Proc. pp. 294 – 297
- [7] R. Wagner, C. Linke, K.-H. Naumann, M. Schnaiter, M. Vragel, M. Gangl, H. Horvath, “A review of optical measurements at the aerosol and cloud chamber AIDA”, *Journal of Quantitative Spectroscopy and Radiative Transfer* **110**, 930 – 949 (2009)
- [8] Z. Ulanowski, C. Stopford, E. Hesse, P. H. Kaye, E. Hirst, and M. Schnaiter, “Characterization of small ice crystals using frequency analysis of azimuthal scattering patterns”, in *10th Int. Conference on Electromagnetic & Light Scattering*; Bodrum, Turkey (2007), Proc. pp. 225 – 228

^a Karlsruhe Institute of Technology
Institute for Meteorology and Climate Research
P.O. Box 3640
76021 Karlsruhe, Germany

^b University of Hertfordshire
Centre for Atmospheric and Instrumentation Research
Hatfield AL10 9AB, United Kingdom

* To whom correspondence should be addressed | Email: martin.schnaiter@kit.edu

Paper presented at the ELS XIII Conference (Taormina, Italy, 2011), held under the APP patronage; published online 15 September 2011.

© 2011 by the Author(s); licensee *Accademia Peloritana dei Pericolanti*, Messina, Italy. This article is an open access article, licensed under a [Creative Commons Attribution 3.0 Unported License](https://creativecommons.org/licenses/by/3.0/).