Corrected Quantum Drift-Diffusion equation via Compressed Chapman-Enskog expansion

Giovanni Frosali and Chiara Manzini Dipartimento di Matematica Applicata "G.Sansone" Università di Firenze - Via S.Marta 3 I-50139 Firenze, Italy

giovanni.frosali@unifi.it, chiara.manzini@unifi.it

One of the most important aspects of quantum theory is the derivation of simple approximate models providing a hydrodynamic description of the transport of charged carriers. Quantum drift-diffusion equations, e.g., are both subject of diverse mathematical studies ([4] and the Refs. therein) and used by engineers in many practical simulations. Our aim is the derivation of a quantum drift-diffusion equation via the modified (compressed) Chapman–Enskog method [3], as the preliminary step towards a rigorous asymptotic analysis of the quantum transport equation, in case a strong external electric field and a relaxation-time collisional operator are included. Therefore, the Wigner equation for the quasi-distribution function w = w(x, v, t) is modified by a BGK operator, that indicates relaxation in a time $1/\nu$ to the equilibrium state w_{eq} . Precisely, the Wigner equation for the quasi-distribution function w = w(x, v, t) is

$$\frac{\partial w}{\partial t} + v \cdot \nabla w - \Theta[V]w = -\nu(w - w_{eq}), \quad (x, v) \in \mathbb{R}^{2d}, \ t > 0,$$

where the potential V enters through the pseudo-differential operator $\Theta[V]$ defined by

$$(\Theta[V]w)(x,v,t) = \frac{i}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} \delta V(x,\eta) \mathcal{F}w(x,\eta,t) e^{iv\cdot\eta} \, d\eta \,,$$

with $\delta V(x,\eta) := \frac{1}{\hbar} \left(V(x + \frac{\hbar\eta}{2m}) - V(x - \frac{\hbar\eta}{2m}) \right)$ and $\mathcal{F}f(\eta)$, which denotes the Fourier transform of w from the variable v to the variable η . We can introduce a small parameter ϵ related to the mean free path, which refers to the relaxation to thermodynamical equilibrium and to the presence of a strong field. Accordingly, in dimensionless form

(0.1)
$$\frac{\partial w}{\partial t} + v \cdot \nabla w - \frac{1}{\epsilon} \Theta[V] w = -\frac{w - w_{eq}}{\epsilon}, \quad (x, v) \in \mathbb{R}^{2d}, \ t > 0.$$

We shall describe the equilibrium state by the $O(\hbar^2)$ -quantum corrected thermodynamical equilibrium function calculated by Wigner in [6] (cf. [2] for an updated discussion). Precisely, we can write the Wigner thermal equilibrium function as

$$w_{eq}(x,v) = n(x,t) \left(F(v) + \hbar^2 F^{(2)}(x,v) \right) + O(\hbar^4), \ F(v) := \left(\frac{\beta m}{2\pi}\right)^{d/2} e^{-\beta m v^2/2},$$

$$F^{(2)}(x,v) := F(v) \left[-\frac{\beta^2}{24m} \sum_{r=1}^d \frac{\partial^2 V}{\partial x_r^2} + \frac{\beta^3}{24} \sum_{r,s=1}^d v_r v_s \frac{\partial^2 V}{\partial x_r x_s} \right],$$

Licensed under the Creative Commons Attribution Noncommercial No Derivatives

G. Frosali et al.

where $n(x,t) = n[w](x,t) := \int w(x,v,t) dv$ is the position density, $\beta \equiv 1/kT$, T is the (constant) temperature and k is the Boltzmann constant (cf. [1]).

We shall consider the Hilbert space $X_k = L^2(\mathbb{R}^{2d}, (1+|v|^{2k})dx dv; \mathbb{R})$ and write (0.1) in the abstract form

(0.2)
$$\epsilon \frac{dw}{dt} = \epsilon Sw + \mathcal{A}w + \mathcal{C}w, \qquad \lim_{t \to 0^+} \|w(t) - w_0\|_{X_k} = 0$$

with

$$\mathcal{A}w := \Theta[V]w, \qquad \mathcal{C}w := -(w - \emptyset w), \quad \Omega w = n(F + \hbar^2 F^{(2)})$$

and the free-streaming operator $Su = -v \cdot \nabla, D(S) = \{u \in X_k \mid Su \in X_k\}.$

The main ingredient of the compressed Chapman-Enskog expansion, as proposed in [3], is the study of the problem (0.2) with $\epsilon = 0$, i.e. of the equation $(\mathcal{A} + \mathcal{C})f = 0$ in the space X_k . We can state that, for any fixed $x \in \mathbb{R}^d$

$$\ker(\mathcal{A}+\mathcal{C}) := \{ cM(v), c \in \mathbb{R} \} \subset X_k^v := L^2(\mathbb{R}^d, (1+|v|^{2k})dv; \mathbb{R})$$

with

(0.3)
$$M(x,v) = \mathcal{F}^{-1} \left\{ \frac{\mathcal{F}F(\eta)}{1 - i\delta V(x,\eta)} \left(1 - \frac{\beta^2 \hbar^2}{24m} \sum_{r,s=1}^d \eta_r \eta_s \frac{\partial^2 V(x)}{\partial x_s x_s} \right) \right\} (x,v).$$

Moreover, for all $h \in X_k^v$, $(\mathcal{A} + \mathcal{C})u = h$ has a solution if and only if $\int h(v) dv = 0$. According to the previous considerations, we can decompose the space X_k as

$$X_k = (X_k)_M \oplus (X_k)^0$$

where $(X_k)_M$ is the eigenspace spanned by M and

$$(X_k)^0 = \left\{ f \in X_k \left| \int f(v) \, dv = 0 \right\}.$$

Let \mathcal{P} be the corresponding spectral projection from (X_k) into $(X_k)_M$, defined by $\mathcal{P}f = M \int f(v) dv$ and $\mathcal{Q} = \mathcal{I} - \mathcal{P}$. Accordingly, we decompose the unknown function $w \in X_k$ as $w = \varphi + \psi = \mathcal{P}w + \mathcal{Q}w$, where φ is called the hydrodynamic part and ψ is called the kinetic part of w. Observe that for all $w \in X_k, \mathcal{P}w = Mn[w]$.

Operating formally on both sides of Eq. (0.2 with the projections \mathcal{P} and \mathcal{Q} , we obtain the following system

$$\begin{cases} \frac{\partial \varphi}{\partial t} = \mathcal{P}S\mathcal{P}\varphi + \mathcal{P}S\mathcal{Q}\psi \\ \frac{\partial \psi}{\partial t} = \mathcal{Q}S\mathcal{P}\varphi + \mathcal{Q}S\mathcal{Q}\psi + \frac{1}{\epsilon}\mathcal{Q}(\mathcal{A} + \mathcal{C})\mathcal{Q}\psi \end{cases}$$

with initial conditions

$$\varphi(0) = \varphi_0 = \mathcal{P}f_0, \quad \psi(0) = \psi_0 = \mathcal{Q}f_0.$$

The compressed asymptotic expansion consists in splitting the unknown functions φ and ψ into the sum of the "bulk" parts $\bar{\varphi}$ and $\bar{\psi}$ and of the initial layer parts $\tilde{\varphi}$ and $\tilde{\psi}$, which

take account of the rapid changes of f for small times $t = \frac{t}{\epsilon}$. Then the bulk hydrodynamic part $\bar{\varphi}$ is left unexpanded and the other parts are expanded in series of ϵ . By disregarding terms of order ϵ^2 , we recover the following equation for the unexpanded function $\bar{\varphi}(x, v, t)$

(0.4)
$$\frac{\partial \bar{\varphi}}{\partial t} = \mathcal{P}S\mathcal{P}\bar{\varphi} - \epsilon \mathcal{P}S\mathcal{Q}(\mathcal{Q}(\mathcal{A} + \mathcal{C})\mathcal{Q})^{-1}\mathcal{Q}S\mathcal{P}\bar{\varphi}$$

 $\bar{\varphi}(x, v, t)$ can still be written as the product $\bar{\varphi}(x, v, t) = M(x, v) n(x, t)$, since we shall consider the contribution of the initial layer part $\tilde{\varphi}$ via an appropriate initial condition. After some algebra manipulations, we write Eq. (0.4) in the following form

(0.5)
$$\frac{\partial n}{\partial t} = -\nabla \cdot \left(n \int v M dv \right) \\ -\epsilon \nabla \cdot \left[\left(\int v \otimes D_2 \, dv \right) \nabla n + n \int v D_1 \, dv \right]$$

where D_2 and D_1 are esplicitly determined by solving appropriate equations containing the operator $\mathcal{Q}(\mathcal{A} + \mathcal{C})\mathcal{Q}$. We remark that the solution of Eq. (0.5) is the correction up to order ϵ^2 of the solution *n* of the continuity equation, in case of strong field. We can calculate the coefficient in the operator \mathcal{PSP} defined above

(0.6)
$$\int v M(x,v) dv = -\nabla V(x),$$

the diffusion tensor D

$$\mathsf{D} := \int v \otimes D_2 \, dv = \frac{\mathcal{I}}{\beta} + \nabla V \otimes \nabla V - \frac{\beta \hbar^2}{12m} (\nabla \otimes \nabla) V + O(\hbar^4) \,,$$

with \mathcal{I} identity tensor, and

$$\int v D_1(x,v) dv = (\nabla \otimes \nabla) V \ \nabla V - \frac{\beta \hbar^2}{12m} \nabla \cdot (\nabla \otimes \nabla) V + O(\hbar^4) \,.$$

Finally, Eq. (0.5) looks like

$$\frac{\partial n}{\partial t} - \nabla \cdot (n \,\nabla V) - \frac{\epsilon}{\beta} \,\Delta \,n - \epsilon \nabla \cdot (\nabla V \otimes \nabla V \,\nabla n) \\ -\epsilon \nabla \cdot \left[n \,(\nabla \otimes \nabla) V \,\nabla V \right] + \frac{\epsilon \beta \hbar^2}{12m} \nabla \cdot \left[(\nabla \otimes \nabla) V \,\nabla n + n \,\nabla \cdot (\nabla \otimes \nabla) V \right] = 0$$

and we can recognize the drift and the diffusion terms, two terms due to the strong field assumption (cf. [5]) and the terms due to the (quantum) Bohm potential (cf. [1]). We shall give a rigorous proof of the results up to now formally derived. In particular, we study the well-posedness for the initial parts of our problem. We shall also prove regularity of the solution of the drift-diffusion equation, starting from sufficiently smooth initial value. Finally we can prove that the asymptotic expansion up to the first order gives an approximation of order ϵ^2 to the solution of the exact problem.

REFERENCES

G. Frosali et al.

- 1. C. Gardner, The quantum hydrodynamic model for semiconductor devices SIAM J.App.Math. 54(2), 1994, 409–427.
- 2. A. Jüngel and D. Matthes, A derivation of the isothermal quantum hydrodynamic equations using entropy minimization, Z. Angew. Math. Mech. 85, 2005, 806–814.
- 3. J.R. Mika and J. Banasiak, Singularly perturbed evolution equations with applications to kinetic theory. Singapore, World Scientific 1995.
- 4. R. Pinnau, A review on the quantum drift diffusion model. *Transp. Theory Stat. Phys.* 31(4-6), 2002, 367–395.
- 5. F. Poupaud, Runaway phenomena and fluid approximation under high fields in semiconductor kinetic theory. Z. Angew. Math. Mech. 72, 1992, 359–372.
- E. Wigner, On the quantum correction for thermodynamic equilibrium, *Phys. Rev.* 40, 1932, 749–759.